

# Filling in the GAP: Occupational Therapy's Role in Interprofessional Geriatric Assessment Team

Ashley Halle, OTD, OTR/L






- 1. Existing program logistics/flow and future directions
- 2. Strategies and mechanisms that led to successful intra-team cohesion and effectiveness
- 3. Opportunities to teach basic principles of geriatric assessment and interprofessional team functioning to students from six health care professions
- 4. Value of Rapid Cycle Quality Improvement (RCQI) to improve processes and experience of care.

# Geriatric Assessment Program (GAP)



- Comprehensive geriatric assessment clinic
- Philosophy
  - Mission To provide beyond exceptional care of older adults with complex health needs through an interprofessional collaborative assessment program.
  - Vision To maximize quality of life for older adults and their families.




Keck School of Medicine of USC



# Why Patients are Referred

- Patients looking for support to age successfully
- Caregivers overwhelmed & needing support
- Connecting with local resources
- Patients are in transitional period
  - Can he/she age in place or does he/she need to transition to a higher level of care?



### The Process

Screening of Referrals Social Work Clinic Visit

Dentist, OT,

Pharmacy, PT,

Physician,

Psychologist

Patient/
Family
Conference
Physician
Assistant
Psychologist












**Home Visit** 

Social Work Occupational Therapy Team
Conference
All team
members



# The Clinic Flow

## Home Evaluations



#### Customized OT Home Evaluation

- Occupational Profile
- Social History & Support

Past Medical History/Past Surgical History
Falls History


Home/Living Situation

**Detailed Environmental Assessment** 

- -- Safety, Accessibility, Adaptive Equipment/DME
- -- Functional mobility & participation in context; meaningful spaces & places



## Clinic Consultation







OT Home Evaluation



OT Clinic Consult

#### OT Rec's

e.g. environmental modifications, AE, changes to routines, alternative occupations, resources, referrals



# Rapid Cycle Quality Improvement (RCQI)

- A quality-improvement method that identifies, implements and measures changes made to improve a process or a system".<sup>5</sup>
- Facilitates improvement through rapid PDSA (Plan, Do, Study, Act) cycles.
- Continually improve health care processes by instituting a constant cycle of innovations or improvements



# RCQI Example

ISSUE	PLAN	DATE	DO	DATE	STUDY	DATE	ACT	DATE
Not all clinicians are credentialed	Work with disciplines and Keck to identify and resolve credentialing procedures.	9 80 80 80	Incorporate substitute clinicians who are credentialed until identified team members are credentialed.	2/17/16	Periodic updates on credential status and HHR training status		Departmental MOUs completed; all clinicians credentialed	4/27/16
Clinic flow is poor due to length of time clinicians work with patients	Install white board and clock	2/17/16	Identify clinician, and ultimately medical assistant, to monitor and remind clinicians.	2/24/16	White board installed, evaluating mechanisms for management and reminders to clinicians	2/24/16	Time allotments added and adjusted. Clinician time in/out logged.	4/13/16



# Case Study – "Jane"





## "Jane" Home Evaluation

#### **BACKGROUND**

- 72 year old African American woman
- Never married, living with brother
- Brother looking for help
- Needs help with her care, specifically concerns related to hygiene and wandering, and wants to optimize her quality of life



# "Jane" - Home Evaluation

- Born in California, 1 of 10 children
- Ivy League education, worked at CNN
- Trauma in 1980 possible victim of sexual assault
  - Attempted suicide by gas in apartment
- Moved back home with mother shortly after
- Parents deceased
- Brother moved back home due to own health issues



# "Jane" - Home Evaluation

- History of small TIAs
- Client is wandering
- Brother stays with her 24/7, receives 52 hours IHSS
- Home in severe disrepair; possible health hazards
- Significant clutter; house not cleaned or maintained for extended period of time
- Brother oversees medications, finances and transportation



# "Jane" - Clinic Consultation

- Stopped caring for self and performing activities around the home
- Extent of hygiene concerns

# "Jane"



### **RESOLUTION**

- Admitted to local gero-psych unit
- Brother working with social work to reconcile options and decide how to proceed

# References



- Anderson, L.A.: Edwards, V.J., Pearson, W.S., et al. (2013). Adult caregivers in the United States: characteristics and differences in well-being, by caregiver age and caregiving status. Prev Chronic Dis 20, E135
- Chrisler J, et al. (2016). Ageism can be hazardous to women's health: Ageism, sexism, and stereotypes of older women in the health care system. J Soc Issues, 72(1), 86–104.
- Chung, J.C., Lai, C.K., Chung, P.M. & French, H.P. (2002) Snoezelen for dementia. Cochrane Database of Systematic Reviews 2002, 4 (CD003152). DOI: 10.1002/14651858.CD003152.
- Counsell, S. R., Callahan, C. M., Clark, D. O., Tu, W., Buttar, A. B., Stump, T. E., & Ricketts, G. D. (2007). Geriatric care management for low-income seniors. JAMA, 298(22), 2623. doi:10.1001/jama.298.22.2623
- Davies, R. (2015). Is Bruges the most dementia-friendly city? The Guardian. 21 April 2015. https://www.theguardian.com/society/2015/apr/21/bruges-most-dementia-friendly-city
- Elder Workforce Alliance (2013). Caring for an aging America: meeting the health care needs of older adults. Retrieved from: https://eldercareworkforce.org/research/issue-briefs/research/ganda/
- Garvey, J., Connolly, D., Boland, F., & Smith, S. M. (2015). OPTIMAL, an occupational therapy led self-management support programme for people with multimorbidity in primary care: A randomized controlled trial. BMC Family Practice, 16(1), . doi:10.1186/s12875-015-0267-0
- Hoffman, S.J., Rosenfield, D., Nasmith, L. (2009) What attracts students to interprofessional education and other health care reform initiatives?. Journal of Allied Health, 38 (3), 75-78.
- Kales, H.C., Gitlin, L.N., Lyketsos, C.G., et al. (2014). Management of neuropsychiatric symptoms of dementia in clinical settings: recommendations from a multidisciplinary expert panel. Journal of the American Geriatrics Society, 62 (4), 1532-5415. http://dx.doi.org/10.1111/jgs.12730
- Kawashima, Ryuta et al. (2015) SAIDO learning as a cognitive intervention for dementia care: a preliminary study. Journal of the American Medical Directors Association, 16 (1), 56 62
- Lookinland S, & Anson K (1995). Perpetuation of ageist attitudes among present and future health care personnel: Implications for elder care. J Adv Nurs, 21(1), 47–56.
- Luo B, Zhou K, Jin E., Newman A, & Liang J (2013). Ageism among college students: A comparative study between US and China. J Cross Cult Gerontol, 28(1), 49-63.
- Giles L et al. (2002). Ageism among health professionals: A comparison of clinical educators and students in physical and occupational therapy. Phys Occup Ther Geriatr, 21(2), 15-26.
- Eskildsen M, & Flacker J (2009). A multimodal aging and dying course for first-year medical students improves knowledge and attitudes. JAGS, 57(8), 1492-1497.
- · Purtilo. R.B. (2005). Ethical dimensions in the health professions (4th ed.) Philadelphia: Elsevier.
- State of California Department of Motor Vehicles Dementia. https://www.dmv.ca.gov/portal/dmv/detail/dl/driversafety/dementia
- State of California Department of Motor Vehicles (2016). Driver Safety Information Medical Conditions and Traffic Safety.
  - https://www.dmv.ca.gov/portal/dmv/?1dmy&urile=wcm:path:/dmv content en/dmv/dl/driversafety/dsmedcontraffic
- Institute for Healthcare Improvement. (2016). IHI triple aim initiative: Better care for individuals, better health for populations, and lower per capita costs. Retrieved from: http://www.ihi.org/engage/initiatives/tripleaim/pages/default.aspx
- Administration on Aging (2014). Future growth, Retrieved from: http://www.aoa.acl.gov/Aging Statistics/Profile/2013/4.aspx
- World Health Organization (WHO). (2010). Framework for action on interprofessional education & collaborative practice. Geneva: World Health Organization. http://whqlibdoc.who.int/hq/2010/WHO HRH HPN 10.3 eng.pdf

## References



- Ahlmen, M., Sullivan, M., & Bjelle, A. (1988). Team versus non-team outpatient care in rheumatoid arthritis. Arthritis and Rheumatism, 31(4), 471-479.
- Boult, C., Boult, L. B., Morishita, L., Dowd, B., Kane, R. L., & Urdangarin, C. F. (2001). A randomized clinical trial of outpatient geriatric evaluation and management. *Journal of the American Geriatrics Society, 49,* 351-359.
- Burns, R., Nichols, L. O., Martindale-Adams, J., & Graney, M. J. (2000). Interdisciplinary geriatric primary care evaluation and management: Two-year outcomes. Journal of the American Geriatrics Society, 48, 8-13.
- Cohen, H. J., Fuessner, J. R., Weinberger, M., Carnes, M., Hamdy, R. C., Hsieh, F., Phibbs, C. & Lavori, P. (2002). A controlled trial of inpatient and outpatient geriatric evaluation and management. New England Journal of Medicine. 346(12), 905-912.
- Counsell, S. R., Callahan, C. M., Clark, D. O., Tu, W., Buttar, A. B., Stump, T. E., & Ricketts, G. D. (2007). Journal of the American Medical Association, 298(22), 2623-2633.
- Donnelly, C., Brenchley, C., Crawford, C., & Letts, L. (2013). The integration of occupational therapy into primary care: a multiple case study design. BMC Family Practice, 14(1). http://dx.doi.org/10.1186/1471-2296-14-60
- Engelhardt, J., Toseland, R. W., Gao, J., & Banks, S. (2006). Long-Term Effects of Outpatient Geriatric Evaluation and Management on Health Care Utilization, Cost, and Survival. Research On Social Work Practice, 16(1), 20-27. http://dx.doi.org/10.1177/1049731505276047
- Kuo, H., Scandrett, K., Dave, J., & Mitchell, S. (2004). The influence of outpatient comprehensive geriatric assessment on survival: a meta-analysis. *Archives Of Gerontology And Geriatrics*, 39(3), 245-254. http://dx.doi.org/10.1016/j.archger.2004.03.009
- Mackenzie, L. & Clemson, L. (2014). Can chronic disease management plans including occupational therapy and physiotherapy services contribute to reducing falls risk in older people?. *Australian Family Physician*, 43(4), 211-215.
- Mackenzie, L., Clemson, L., & Roberts, C. (2013). Occupational therapists partnering with general practitioners to prevent falls: Seizing opportunities in primary health care. *Australian Occupational Therapy Journal*, 60(1), 66-70. http://dx.doi.org/10.1111/1440-1630.12030
- McColl, M., Shortt, S., Godwin, M., Smith, K., Rowe, K., O'Brien, P., & Donnelly, C. (2009). Models for Integrating Rehabilitation and Primary Care: A Scoping Study. Archives Of Physical Medicine And Rehabilitation, 90(9), 1523-1531. http://dx.doi.org/10.1016/j.apmr.2009.03.017
- Reuben, D. B., Frank, J. C., Hirsch, S. H., McGuigan, K. A., & Maly, R. C. (1999). A randomized clinical trial of outpatient comprehensive geriatric assessment coupled with an intervention to increase adherence to recommendations. *Journal of the American Geriatrics Society*, 47, 269-276.
- Stuck, A. E., Siu, A. L., Weiland, G. D., Adams, J., & Rubenstein, L. Z. (1993). Comprehensive geriatric assessment: a meta-analysis of controlled trials. Lancet, 342, 1032-36.