





Background

- Setting up a new service.
- Measuring outcomes has been a focus of healthcare in UK.
- Anecdotal experience.
- Final year of MSc.

(Department of Health 2010, College of Occupational Therapists 2013, Richards et al 2015).





The Study

Aim: To discover how occupational therapists in oncology and palliative care are evaluating their practice.

- Study Design
- Sample
- Data analysis

ARDIFF Small sample sizes More training needed in use UNIVERSITY of outcome measures PRIFYSGOL Difficulties recruiting patients More work needed Little known about OT role with people with long term conditions Lack of research High attrition rates Developing an outcome measure for all palliative stages = challenge

(Norris 1999, Kealey and McIntyre 2005, Harrison – Paul and Drummond 2006, Pearson et al 2007, Stevens et al 2010, Hammill et al 2014, Phipps and Cooper 2014)









- Limited use of standardised outcome measures.
- Reliance on patient evaluation tools:
 8 out of 10 participants using patient evaluation questionnaires only.
- Lack of training in use of outcome measures:
 7 out of 10 participants had training but this was out of date and not specific.



Recommendations



• Training for students and therapists re: evaluating practice.

Support from services.

• More research is needed to highlight the contribution of occupational therapy.



Royal College of Occupational Therapists
Specialist Section

Oncology & Palliative Care

Publication

Research



Challenges are opportunities in disguise...

CowpeL1@cardiff.ac.uk